

In Vitro Antimicrobial Resistance of Urinary *Escherichia coli* Isolates among U.S. Outpatients from 2000 to 2010

Guillermo V. Sanchez,^a Ronald N. Master,^b James A. Karlowsky,^c and Jose M. Bordon^d

School of Public Health and Health Services, George Washington University, Washington, DC, USA^a; Quest Diagnostics Nichols Institute, Chantilly, Virginia, USA^b; Department of Medical Microbiology and Infectious Diseases, University of Manitoba, Winnipeg, Manitoba, Canada^c; and Section of Infectious Disease, Providence Hospital, Washington, DC, USA^d

This study examines *in vitro* antimicrobial resistance data from *Escherichia coli* isolates obtained from urine samples of U.S. outpatients between 2000 and 2010 using The Surveillance Network (TSN). Antimicrobial susceptibility results ($n = 12,253,679$) showed the greatest increases in *E. coli* resistance from 2000 to 2010 for ciprofloxacin (3% to 17.1%) and trimethoprim-sulfamethoxazole (TMP-SMX) (17.9% to 24.2%), whereas nitrofurantoin (0.8% to 1.6%) and ceftriaxone (0.2% to 2.3%) showed minimal change. From 2000 to 2010, the antimicrobial resistance of urinary *E. coli* isolates to ciprofloxacin and TMP-SMX among outpatients increased substantially.

Antimicrobial resistance significantly increases patient morbidity, costs of treatment, rates of hospitalization, and use of broad-spectrum agents (7). Resistant *Escherichia coli* isolates are associated with decreases in clinical cure rates and higher risk of recurrence (17, 20).

Several studies have described the *in vitro* susceptibility of *E. coli* isolates among outpatients in the United States, and most of these studies have focused on women. The most-recent published data available for U.S. outpatients were collected from April 2003 to June 2004 and suggested the levels of antimicrobial resistance of urinary *E. coli* isolates to be 39.3% for ampicillin, 22.6% for trimethoprim-sulfamethoxazole (TMP-SMX), 6.8% for ciprofloxacin, and 1.4% for nitrofurantoin (22). Other, smaller regional studies suggested a continued trend of rising resistance in the outpatient setting (8, 13, 16).

Limited data are available to describe long-term trends in antimicrobial resistance of *E. coli* isolates among outpatients in the United States. The objective of this study was to examine trends of antimicrobial resistance of urinary *E. coli* isolates among outpatients in the United States from 2000 to 2010.

Antimicrobial susceptibility test results were obtained from The Surveillance Network (TSN) Database—USA (Eurofins Medinet, Chantilly, VA). This surveillance database collects data from over 200 institutions in the United States, and antimicrobial susceptibility testing is performed on-site by each laboratory in

accordance with FDA-approved testing methods and interpreted using Clinical and Laboratory Standards Institute (CLSI)-recommended breakpoints. TSN data have been used before to evaluate trends in antimicrobial resistance, and further details of quality control have been described previously (5, 9, 18, 19).

The present study included antimicrobial susceptibility data for urinary *E. coli* isolates obtained from U.S. outpatients between 2000 and 2010. Outpatients are defined as individuals who visited emergency departments, hospital-based outpatient clinics, and physicians' offices. *E. coli* isolates with intermediate susceptibility were not classified as being resistant. The outcomes of interest in this study were the changes in and most-recent prevalence of antimicrobial resistance to commonly prescribed oral agents used to treat urinary tract infections (UTIs). Cephalothin and ceftriaxone, both intravenously administered agents, were selected as narrow-

Received 7 November 2011 Returned for modification 6 December 2011

Accepted 3 January 2012

Published ahead of print 17 January 2012

Address correspondence to Jose M. Bordon, jbordon@provhosp.org.

Copyright © 2012, American Society for Microbiology. All Rights Reserved.

doi:10.1128/AAC.06060-11

TABLE 1 Annual rates of resistance of urinary *Escherichia coli* isolates to select antimicrobials among all outpatients from 2000 to 2010^a

Antimicrobial agent	No. of test results	Antimicrobial resistance rate (%) for indicated yr											Total change (%) from 2000–2010 ^b
		2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	
Ciprofloxacin	1,836,598	3	3.7	5	6.2	7.5	9.7	12.2	14.1	16	16.3	17.1	14.1
TMP-SMX	2,034,254	17.9	17.8	18.1	18.2	19.4	20.2	20.8	21.9	22.9	23	24.2	6.3
Ampicillin	2,002,221	38.2	38.1	38.2	38.2	38.2	39.4	40.5	41.5	42.4	42.4	43.4	5.2
Cefuroxime	806,659	1.5	1.9	2.1	2.3	2.4	2.3	2.9	3.2	3.7	4	5	4.5
Cephalothin	502,231	14	15.8	14.6	15	14.5	15.7	16.9	16.6	16.4	16	18.1	4.1
Tetracycline	580,328	22.6	22.1	22.1	21.1	21.9	22	23.5	23.7	24.4	24.5	24.9	2.3
Ceftriaxone	1,759,006	0.2	0.2	0.3	0.3	0.5	0.6	0.9	1.2	1.6	1.9	2.3	2.1
Nitrofurantoin	1,972,633	0.8	1	1.1	1.1	1	1.1	1.4	1.5	1.5	1.6	1.6	0.8
Amox-Clav	759,749	5	4.1	5.3	4.5	3.7	3.8	5.6	8.2	9.9	6.6	5.3	0.3

^a Isolates demonstrating intermediate susceptibility were not counted as resistant. Amox-Clav, amoxicillin-clavulanate; TMP-SMX, trimethoprim-sulfamethoxazole.

^b All antimicrobial agents demonstrated statistically significant changes from 2000 to 2010.




FIG 1 Cumulative annual change in *E. coli* antimicrobial resistance in outpatient urinary *E. coli* isolates from 2001 to 2010. Amox/Clav, amoxicillin-clavulanate.

and broad-spectrum cephalosporin surrogates, respectively, for their orally administered formulations.

A chi-square test was performed for each antimicrobial agent to determine whether a significant difference existed between resistance rates observed in 2000 and those observed in 2010. An alpha level of 0.05 was used. Analyses were performed using Statistical Analysis Software (SAS) version 9.1.

Antimicrobial susceptibility results ($n = 12,253,679$) for urinary *E. coli* isolates obtained from outpatients in the United States from 2000 to 2010 were examined (Table 1). The greatest increases in resistance among isolates obtained from all outpatients from 2000 to 2010 were observed for ciprofloxacin (from 3% in 2000 to 17.1% in 2010) and TMP-SMX (17.9% to 24.2%) (Fig. 1). Conversely, over the same time period, nitrofurantoin (0.8% to 1.6%), amoxicillin-clavulanate (5% to 5.3%), and ceftriaxone (0.2% to 2.3%) demonstrated only small changes in resistance. In 2010, ampicillin, tetracycline, cephalothin, and cefuroxime showed antimicrobial resistance rates of 43.4%, 24.9%, 18.1%, and 5.0%, respectively.

Emerging antimicrobial resistance of *E. coli* in the outpatient setting is well documented (2, 3, 7, 12, 14). In the early 2000s, quinolones surpassed sulfa drugs as the most common class of antimicrobials prescribed by clinicians to treat uncomplicated UTIs (10). This increase in provider use of fluoroquinolones may account for the rapid rise in antimicrobial resistance of *E. coli* to ciprofloxacin, as resistance to this agent has been shown to correlate with the level of use (8, 15, 21). Due to the propensity of *E. coli* to acquire resistance to this agent, use of ciprofloxacin for empirical treatment of UTIs in outpatients should be used sparingly and only where local antimicrobial resistance rates remain low (4).

Antimicrobial resistance of urinary *E. coli* isolates to TMP-SMX continued to increase from 2000 to 2010, a trend that has continued for decades (1, 6, 11). In the 2010 IDSA guidelines for treating acute uncomplicated cystitis in women, TMP-SMX is recommended as the second-line antimicrobial agent (4). Our data are consistent with previous reports regarding increases in antimicrobial resistance of urinary *E. coli* isolates to TMP-SMX and potential subsequent decreases in its efficacy as empirical therapy among U.S. outpatients.

Levels of antimicrobial resistance of *E. coli* to cephalothin, the narrow-spectrum oral cephalosporin surrogate, were higher than those for expanded-spectrum (cefuroxime) and broad-spectrum (ceftriaxone) cephalosporins. It is important to note that while the

absolute change in antimicrobial resistance to ceftriaxone was small (0.2% in 2000 to 2.3% in 2010), there was a 10-fold increase in resistance of *E. coli* to this agent over the study time period. Though these changes do not bear immediate clinical significance, future surveillance of antimicrobial resistance to this agent is warranted.

The *in vitro* antimicrobial resistance rates among *E. coli* isolates in our investigation were consistent with those reported previously (6, 11, 22). For example, antimicrobial resistance of *E. coli* to nitrofurantoin demonstrated little change over our study time period, a finding that is consistent with the resistance prevalence reported in the NAUTICA study (22).

It is important to note the strengths and limitations of our data. The strengths of our study include the large number of isolates, the variety of antimicrobial agents studied, the large number of reporting institutions within the United States, the long time period for which data were reported, and the geographically representative distribution of isolates from TSN. The limitations of our study include a lack of central laboratory testing, the use of multiple susceptibility test methods, and an assumed underrepresentation of isolates from those for whom empirical treatment was successful. These data should be interpreted with caution. Although traditional *in vitro* surveillance systems are well designed to provide insight into overall trends and prevalence of antimicrobial resistance, they are not meant to guide antimicrobial therapy in the management of individual clinical cases.

In summary, our study shows that from 2000 to 2010, antimicrobial resistance of urinary *E. coli* isolates to ciprofloxacin and TMP-SMX increased substantially but that resistance to nitrofurantoin and ceftriaxone remained low. Given the frequency with which UTIs are treated empirically, compounded with the speed that *E. coli* acquires resistance, prudent use of antimicrobial agents remains crucial.

REFERENCES

- Diekema DJ, et al. 2004. Antimicrobial resistance trends and outbreak frequency in United States hospitals. *Clin. Infect. Dis.* 38:78–85.
- Gordon KA, Jones RN. 2003. Susceptibility patterns of orally administered antimicrobials among urinary tract infection pathogens from hospitalized patients in North America: comparison report to Europe and Latin America. Results from the SENTRY Antimicrobial Surveillance Program (2000). *Diagn. Microbiol. Infect. Dis.* 45:295–301.
- Gupta K, Hooton TM, Stamm WE. 2001. Increasing antimicrobial resistance and the management of uncomplicated community-acquired urinary tract infections. *Ann. Intern. Med.* 135:41–50.

4. Gupta K, et al. 2011. International clinical practice guidelines for the treatment of acute uncomplicated cystitis and pyelonephritis in women: a 2010 update by the Infectious Diseases Society of America and the European Society for Microbiology and Infectious Diseases. *Clin. Infect. Dis.* 52:e103–e120.
5. Gupta K, Sahm DF, Mayfield D, Stamm WE. 2001. Antimicrobial resistance among uropathogens that cause community-acquired urinary tract infections in women: a nationwide analysis. *Clin. Infect. Dis.* 33:89–94.
6. Hooton TM. 2003. The current management strategies for community-acquired urinary tract infection. *Infect. Dis. Clin. North Am.* 17:303–332.
7. Hooton TM, Besser R, Foxman B, Fritsche TR, Nicolle LE. 2004. Acute uncomplicated cystitis in an era of increasing antibiotic resistance: a proposed approach to empirical therapy. *Clin. Infect. Dis.* 39:75–80.
8. Johnson L, et al. 2008. Emergence of fluoroquinolone resistance in outpatient urinary *Escherichia coli* isolates. *Am. J. Med.* 121:876–884.
9. Jones ME, et al. 2004. Antibiotic susceptibility of bacteria most commonly isolated from bone related infections: the role of cephalosporins in antimicrobial therapy. *Int. J. Antimicrob. Agents* 23:240–246.
10. Kallen AJ, Welch HG, Sirovich BE. 2006. Current antibiotic therapy for isolated urinary tract infections in women. *Arch. Intern. Med.* 166:635–639.
11. Karlowsky JA, Kelly LJ, Thornsberry C, Jones ME, Sahm DF. 2002. Trends in antimicrobial resistance among urinary tract infection isolates of *Escherichia coli* from female outpatients in the United States. *Antimicrob. Agents Chemother.* 46:2540–2545.
12. Kashanian J, et al. 2008. Nitrofurantoin: the return of an old friend in the wake of growing resistance. *BJU Int.* 102:1634–1637.
13. Khawcharoenporn T, Vasoo S, Ward E, Singh K. 2010. High rates of quinolone resistance among urinary tract infections in the ED. *Am. J. Emerg. Med.* [Epub ahead of print.] doi:10.1016/j.ajem.2010.09.030.
14. Lautenbach E, et al. 2004. Longitudinal trends in fluoroquinolone resistance among Enterobacteriaceae isolates from inpatients and outpatients, 1989–2000: differences in the emergence and epidemiology of resistance across organisms. *Clin. Infect. Dis.* 38:655–662.
15. MacDougall C, Powell JP, Johnson CK, Edmond MB, Polk RE. 2005. Hospital and community fluoroquinolone use and resistance in *Staphylococcus aureus* and *Escherichia coli* in 17 US hospitals. *Clin. Infect. Dis.* 41:435–440.
16. Olson RP, Harrell LJ, Kaye KS. 2009. Antibiotic resistance in urinary isolates of *Escherichia coli* from college women with urinary tract infections. *Antimicrob. Agents Chemother.* 53:1285–1286.
17. Raz R, et al. 2002. Empiric use of trimethoprim-sulfamethoxazole (TMP-SMX) in the treatment of women with uncomplicated urinary tract infections, in a geographical area with a high prevalence of TMP-SMX-resistant uropathogens. *Clin. Infect. Dis.* 34:1165–1169.
18. Sahm DF, Marsilio MK, Piazza G. 1999. Antimicrobial resistance in key bloodstream bacterial isolates: electronic surveillance with the Surveillance Network Database—U. S. A. *Clin. Infect. Dis.* 29:259–263.
19. Sahm DF, Brown NP, Yee YC, Evangelista AT. 2008. Stratified analysis of multidrug-resistant *Escherichia coli* in US health care institutions. *Postgrad. Med.* 120:53–59.
20. Talan DA, et al. 2000. Comparison of ciprofloxacin (7 days) and trimethoprim-sulfamethoxazole (14 days) for acute uncomplicated pyelonephritis pyelonephritis in women: a randomized trial. *JAMA* 283:1583–1590.
21. Zervos MJ, et al. 2003. Relationship between fluoroquinolone use and changes in susceptibility to fluoroquinolones of selected pathogens in 10 United States teaching hospitals, 1991–2000. *Clin. Infect. Dis.* 37:1643–1648.
22. Zhanel GG, et al. 2006. Antibiotic resistance in *Escherichia coli* outpatient urinary isolates: final results from the North American Urinary Tract Infection Collaborative Alliance (NAUTICA). *Int. J. Antimicrob. Agents* 27:468–475.